M04B MILK WITHDRAWAL PERIODS FOR DRUGS COMMONLY USED BY DAIRY FARMERS SUPPLYING BAMAKASI

Below are withdrawal periods of chemicals and medicines commonly found in **Bamakasi**. This information is taken from data sheets and is to the best of ZDTP's knowledge accurate as at 13 December 2018. As information may change from time to time, **PLEASE** always check the data sheet / label of the medicine or chemical being given and if you have any questions, contact your local Veterinary Officer.

Type of Drug	Brand Name	Uses	Active ingredients	Milk Withdrawal Period
Antibiotic	Mega pen	Treatment of mastitis, coughing and after calving	Procaine Penicillin G and Benzathine Penicillin	2 days (48 hours)
	Penstrep 400	Treatment of mastitis	Benzathine penicillin	2.5 days (60 hours)
	Oxytetracycline 20% LA	Treatment of Salmonellae, Leptospira, Mycoplasmas, Rickettsia and Chlamydia	Oxytetracycline 200mg	7 days
Parasite treatment (ticks, words, blood parasites)	Imochem-120	Prophylactic treatment of Babesiosis and Anaplasmosis	Imidocarb dipropionate	21 days
	Supermec	Deworming	Ivermectin	0 days
	Miltraz	Dipping	Ecotraz	0 days
	Tick rid	Dipping	Tritax	
Anti - inflammatory	Dexaject	Treatment of udder inflammation	Dexamethasone	3 days (72 hours)
Vitamin	Vitamin AD3E	Vitamin supplement	Vitamin A, D, E	0 days